

RESEARCH

Open Access



The role of co-infections and hormonal contraceptives in cervical intraepithelial neoplasia prevalence among women referred to a tertiary hospital in Western Kenya

Calleb George Onyango^{1*}, Lilian Ogonda¹ and Bernard Guyah¹

Abstract

Background Screening for co-infections with HIV, HSV-2 and *Chlamydia trachomatis* (CT) among high-risk human papilloma virus (hr-HPV) positive women, coupled with enhanced counseling on contraceptives use remains essential in alleviating high morbidity of cervical cancer (CC). The aim of this study was to determine the prevalence of cervical intraepithelial neoplasia (CIN) among women referred for CC screening at a referral hospital in Kisumu County, Kenya; and to establish the role of co-infection and hormonal contraceptives on CIN.

Method In a cross-sectional study, we collected HPV, HIV, HSV-2 and CT data, cervical cytology results, and demographic information from 517 referrals. Blood samples were obtained for HIV and HSV-2 tests; urine for CT test, cervical swabs for hr-HPV test and colposcopic biopsy for histology confirmation after visual inspection with acetic acid (VIA).

Results The overall prevalence of CIN was 18.4% (95/517) with CIN1 observed in 56(29.6%), CIN2 in 27(14.3%), CIN3 and above (CIN3+) in 12(6.3%) and normal biopsy in 94(49.7%) of the patients out of which high grade CIN2 and above (CIN2+) was 7.54% (39/517) equivalent to 32.5 per 100,000 women per year. In a univariate analysis; HPV/HIV co-infection (infected vs. uninfected: OR 2.79; 95% CI 1.56–5.10, $p < 0.001$); HPV/HSV-2 co-infection (infected vs. uninfected: OR 2.41; 95% CI: 1.12–5.46, $p < 0.024$); HPV/CT co-infection (infected vs. uninfected: OR 3.83; 95% CI 1.84–8.51, $p < 0.001$) were found to be significantly associated with CIN. Additionally, hormone-containing intra uterine device (HIUD) contraceptives (users vs. none users: OR 1.43; 95% CI 0.28–10.9, $p < 0.017$) were also associated with CIN.

Conclusion Co-infections with HIV, HSV-2 or *Chlamydia trachomatis* and use of HIUD were associated with increased risk of testing positive for CIN in HPV positive women. Although the overall prevalence of CIN was high, high-grade CIN2+ was comparable to the rates reported earlier. Therefore, population screening for co-infections alongside hr-HPV is desirable and is likely to reduce the burden of CIN in the region. Besides, women positive for hr-HPV and opting for contraceptives ought to be counseled about the possible positive and negative side-effects of different contraception options.

*Correspondence:
Calleb George Onyango
onyangocalleb49@gmail.com

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

Keywords Co-infections, Hormonal contraceptives, HPV, HSV-2, HIV, CT, CIN, Kenya

Introduction

Cervical intraepithelial neoplasia (CIN), the precursor of cervical cancer remain the greatest threat to the reproductive health of many women especially those living in low income, high HIV burden countries in sub Saharan Africa [1]. Globally, the disease is ranked 4th in both incidence and cancer-related mortality amongst women with an estimated 660,000 new cases and 350,000 deaths annually, and accounts for about 13.1% of all new female cancers globally [2]. However, Kenya among other East African countries remains the most affected with an estimated age-standardized incidence and mortality rates of 40.1 and 30.0 per 100,000 respectively [3].

Despite improved preventive strategies including public mobilization for early screening, reduced exposure to infectious agents through community awareness campaign, enhanced immunization to targeted age group, and a concerted campaign to reduce risky behaviors that promotes the spread of the disease [4], Kenya has continued to experience higher incidence and mortality of 5236 and 3211 respectively, annually [1] with prevalence in Kisumu estimated at 8.2% [5]. Besides, infection with HIV confers susceptibility to human papilloma virus (HPV), the causative agent of cervical cancer; and women infected with HIV are at high risk of contracting the disease [6]. Women infected with HIV experience a decline in both the number and function of CD4⁺ T cells leading to high rate of contracting HPV and reduction of chance for spontaneous clearance [7]. Although Kisumu continue to register high burden of HIV [8], data on the prevalence of CIN in the region remain scanty.

More importantly, high risk human papilloma virus (hr-HPV), particularly HPV types 16 and 18, remain the major causative agent of most malignant and premalignant lesions of the cervix, presenting nearly 99% cases of cervical cancer [4]. The virus penetrates the basal layers of epithelial cells through micro-abrasion of the transformation zone of the cervix. Once inside the cell, the viral genome can express E6 and E7 oncogenes which play critical roles in the transformation of infected cells by inhibiting the cell cycle regulating proteins such as p53 and retinoblastoma gene product (pRB), respectively [9]. This inhibition leads to uncontrolled cell proliferation and cervical dyskaryosis following a series of proteolytic degradations [9]. In Kenya, about 9.1% women in the general population are estimated to harbor cervical hr-HPV types 16 /18 infection which contribute to more than 63% of all invasive cervical cancers in the Country [1]. Additionally, higher prevalence of 9.9% have been reported among HIV-infected women in Kisumu [10] suggesting the likelihood of higher burden of CIN in the region.

Moreover, co-infections with herpes simplex virus type 2 (HSV-2) serve as an independent predictor for cervical cancer owing to its role in facilitating HIV acquisition and transmission among sexually active community [11]. Precisely, the virus evolved strategies that counteract caspase activation and apoptosis by encoding anti-apoptotic viral proteins such as ribonucleotide reductase large subunit (R1) [12] leading to the persistence of HPV within the cervix. Although high prevalence of HIV / HSV-2 co-infection have been reported among the local fishing community [13], the role of HSV-2 / HPV co-infection in CIN remain scanty. Equally, *Chlamydia trachomatis* (CT) like other intracellular pathogens have been shown to possess the potential of altering gene expression and protein production in cervical basal cells leading to induction of host genome duplicate that results in aneuploidy and chromosome instability [14]. As *Chlamydia trachomatis* induced DNA double-strand breaks, it simultaneously inhibits proper DNA damage response and repair mechanisms rendering host cells prone to loss of genetic integrity and transformation [15]. Although higher prevalence of HIV / *Chlamydia trachomatis* co-infection have been reported in the region [16] the role of co-infections on CIN prevalence in the region is scanty.

Elsewhere, studies have shown that hormonal contraceptives use increase the risks of developing cervical cancer especially when used for extended period [17]. Specifically, the upstream regulatory region (URR) of HPV 16 genome that mediate transcriptional control of HPV genes contains promoter elements that are activated by persistent interaction with steroid hormones [18]. These hormones binds to specific glucocorticoid-response receptors within the HPV, enhancing the expression of E6 and E7. In turn E6 and E7 bind to and degrade p53 gene product, leading to a apoptotic failure and the development of CIN [18]. Locally, adolescent girls and young women form the majority of contraceptives users with higher preference being injectable and implants [19]. We examined the influence of hormonal contraceptives on CIN prevalence in the region.

Method

Study design

In a cross-sectional study, 517 women aged 25 to 65 years referred to Jaramogi Oginga Odinga Teaching and Referral Hospital (JOOTRH) from peripheral facilities with vaginal or cervix abnormalities between the years 2021 to 2023 were consecutively enrolled in the study after excluding 112 due to insufficient cytological samples (Fig. 1). Eligibility included: (1) women with a history of sexual activity, (2) initial visual inspection with acetic acid




Fig. 1 Flowchart of participants in the study

positive result from referral facilities and (3) informed consent. The exclusion criteria included women: (1) who were pregnant; (2) who had vaginal medication 2 days prior to the screening day (3) those with hysterectomy, muco-purulent discharge, active vaginal bleeding. Nurses in the cervical screening clinic identified potential participants who attended the clinic and explained to them details of the study. Written informed consent was obtained in Kiswahili or English before study enrollment. Clinical examination involved a gynecological examination with inspection of the cervix uteri and collection of specimens by a gynecologist in a separate room. Cervical sample were collected for HPV DNA testing in Preserv-Cyt® Solution (Hologic) using the Cervex-Brush® (Rover). The sample was then stored at ambient temperature until tested. Finally, the cervix was examined after the application of 5% acetic acid (VIA), and biopsy obtained from cervixes that expressed abnormalities suggestive of neoplasia and recorded in datasheet. All women were informed about the VIA result immediately, but for laboratory results, they were booked to collect after one month in the next clinic review visit.

Laboratory testing

HIV infection was tested using Determine HIV-1/2 and confirmed with First Response HIV1/2 card tests; HSV-2 was tested using HerpeSelect-2 enzyme immunoassay; HPV was tested using Gene Xpert HPV assay (Cepheid, Sunnyvale, California, United States [US]). Gene Xpert HPV is based on a multiplex real-time PCR targeting E6 and E7 oncogenes of 14 h-HPV genotypes. The amplification was performed in five fluorescent channels that identifies five groups namely; HPV16, HPV18/45, HPV31/33/35/52/58, HPV51/59, and HPV39/56/66/68, and the results interpreted using GeneXpert software version 4.8 (Cepheid). *Chlamydia trachomatis* was tested using Gene Xpert CT/NG assay (Cepheid, Sunnyvale, California, United States [US]) with primers and probes for the detection of specific chromosomal sequence (serovariants D-K) expressed by the ompA gene of the bacteria in urine sample; while biopsy was tested using hematoxylin and eosin stains.

Data collection

As enrollment continues, information was collected from participants on socio-demographic status and relevant sexual and reproductive health issues including HIV and ART status, contraceptive use, types and duration and

the number of children. Study information was collected electronically except consent forms, a copy of which was issued to the participants after signing. The consent forms with participants' signatures and national ID numbers were collected and secured in locked file cabinets. Labels of laboratory sample were handwritten and contained a computer-generated subject identifier and sample date.

Statistical analysis

Continuous variables were summarized using mean and standard deviation. Categorical variables were summarized using percentages. Prevalence of CIN, HPV, HIV and HSV-2 positivity were calculated overall. A chi-square test was used to compare proportions within group and logistic regression to estimate associations between demographic characteristics and co-infections and the prevalence of CIN.

Ethics considerations

The study approval was obtained from the local review board at Maseno University and JOOTRH Kisumu, Kenya.

Results

Of the 517 eligible participants, the mean age standard deviation was 34.65 ± 7.45 , age range 22–57 years with a median (IQR) of 33 (28, 39) of whom majority 456(88%) were married with children ranging 3 to 4, and implant (34.4%) and injectable (39.3%) were the most preferred contraceptives with a duration median (IQR) of 11 (7, 5) years (Table 1). Among the participants examined, 123(24%), 63(12%), 48(8.9%) and 410(79%), tested positive for HIV, CT, HSV-2 and hr-HPV infections respectively. All colposcopy suggestive of CIN were further subjected to colposcopically guided biopsy for histology examination. Altogether, 189 biopsy specimens were processed and examined, out of which CIN1 were 56(29.6%); CIN2 were 27(14.3%); CIN3+ were 12(6.3%); while normal cervix were 94(49.7%). Overall, the prevalence of CIN was 18.4% (95/517) of which high grade CIN2 and

Table 1 Demographic and clinical characteristics of study participants

Variable	Overall, N = 517 ¹	Histological diagnosis				
		CIN1, N = 56 ¹	CIN2, N = 27 ¹	CIN3+, N = 12 ¹	normal, N = 94 ¹	unavailable, N = 328 ¹
Age	33 (28, 39)	38(34,46)	46(39,48)	43(39,4)	37(32,42)	30 (27, 36)
Marital status						
Married	456 (88%)	46 (82%)	18 (67%)	10 (83%)	89 (95%)	293 (89%)
Single	35 (6.8%)	1 (1.8%)	0 (0%)	1(8.3%)	3 (3.2%)	30 (9.1%)
Widow	26 (5.0%)	9 (16%)	9 (33%)	1(8.3%)	2 (2.1%)	5 (1.5%)
Parity	3.0(3.0,4.0)	4.0(3.0,4)	4.0(4.0,5)	4.0(3.0,5.0)	4.0(3.0,5)	3.0(3.0, 4.0)
Contraceptive's type						
Hormonal-IUD	29 (5.6%)	4 (7.1%)	3 (11%)	0 (0%)	2 (2.1%)	20 (6.1%)
Implant Jadele / Norplant	179 (35%)	24 (43%)	8 (30%)	3 (25%)	36 (38%)	108 (33%)
injectable /DepProvera	202 (39%)	18 (32%)	6 (22%)	5 (42%)	43 (46%)	130 (40%)
None	50 (9.7%)	9 (16%)	10 (37%)	3 (25%)	9 (9.6%)	19 (5.8%)
Oral pill	57 (11%)	1 (1.8%)	0 (0%)	1 (8.3%)	4 (4.3%)	51 (16%)
Duration (Yrs.)	11 (7, 15)	13(11,17)	13 (0, 17)	15 (8,18)	13(10, 19)	10 (7, 13)
VIA						
negative	341 (66%)	35 (63%)	11(41%)	5 (42%)	58 (62%)	232 (71%)
positive	176 (34%)	21 (37%)	16(59%)	7 (58%)	36 (38%)	96 (29%)
HPV test						
negative	107 (21%)	3 (5.4%)	2 (7.4%)	1 (8.3%)	12(12.8)	89 (27%)
positive	410 (79%)	53(94.6)	25(92.6)	11(91.7%)	82 (87%)	239 (73%)
HIV status						
negative	394 (76%)	22 (39%)	12(44%)	6 (50%)	63 (67%)	291 (89%)
positive	123 (24%)	34 (61%)	15(56%)	6 (50%)	31 (33%)	37 (11%)
HSV-2						
negative	471(91%)	40(71%)	23(85%)	9(75%)	83(88%)	316(96%)
positive	46(8.9%)	16(29%)	4(15%)	3(25%)	11(12%)	12(3.7%)
Chlamydia						
Negative	454 (88%)	35 (63%)	20(74%)	8 (67%)	83 (88%)	308 (94%)
Positive	63 (12%)	21 (38%)	7 (26%)	4 (33%)	11 (12%)	20 (6.1%)

CIN1, CIN2, CIN3: Cervical intraepithelial neoplasia 1, 2 or 3, HSV2: herpes simplex virus 2, ¹Median (IQR); n (%)

above (CIN2+) was 7.54% (39/517) equivalent to 32.5 per 100,000 women per year. Of the 95 women positive for CIN lesions, hr-HPV subgroup 16 and subgroup 18/45 were the most common (Fig. 2).

Correlates associated with CIN positivity

In a Univariate analysis (adjusting for age), the study found widows (OR 11.4; 95% CI 3.18–73.2, $p < 0.001$) were more than 10 times likely to test positive for CIN, while single women [OR 0.80; 95% CI 0.10–4.96, $p = 0.001$] were less likely to test positive for CIN as compared to their married counterparts. Women co-infected with HIV (OR = 2.79; 95% CI 1.56–5.10, $p < 0.001$); were two times more likely to test positive for CIN as compared to HIV uninfected counter parts. Additionally, women co-infected with HSV-2 (OR = 2.41, 95% CI: 1.12–5.46, $p < 0.024$); were also twice more likely to test positive for CIN as compared to HSV-2 uninfected counter parts. Women co-infected with *Chlamydia trachomatis* (OR = 3.83; 95% CI 1.84–8.51, $p < 0.001$); were three times more likely to test positive for CIN as compared to uninfected counter parts. Moreover, women using HIUD contraceptives (OR 1.43; 95% CI 0.28–10.9, $p < 0.017$) were more likely to test positive for CIN compared to none users (Table 2).

Discussion

The present study was set to determine the prevalence and determinants of CIN among women in Kisumu County, Kenya. Although continuous monitoring and reporting on prevalence of CIN is essential for estimating the risk of developing cervical cancer and optimizing screening strategy for early detections and treatment [4] this is the first study to comprehensively examine the

carriage rate of CIN exclusively in HIV endemic region of western Kenya.

Findings of the study revealed that the overall prevalence of cervical intraepithelial neoplasia (CIN) was 18.4% (95/517) of which high grade CIN2 and above (CIN2+) was 7.54% (39/517) equitable to approximately 32.5 per 100,000 women per year; relatively lower than the national incidence of 40.1 per 100, 000 women per year [3], as well as the neighbourhood Uganda (56.2 per 100,000 women per year) [20] and Tanzania (54.0 per 100, 000 women per year) [21]. Although previous estimate had reported higher prevalence of 27.9% severe dysplasia in the same region [22] as well as 21.4% prevalence in Nairobi County [23], our finding was within the range of 3.7 – 22.6% reported around Africa [24] and comparable to 8.2% prevalence report earlier by Mungo et al. in the same region [5] suggesting the disease could be widespread locally. This finding is a reflection of high HPV prevalence of 9.1% in the general population [1] as well as 51.1% prevalence among sexually active men of Kisumu County [25] majorly attributed to HPV subtypes 16 and 18 [26]. In this study, hr-HPV subgroup 16 and 18/45 were the most prevalent and major cause of CIN in the region contributing to nearly half of all new cases detected. Higher prevalence of HPV genotype 16 have been reported earlier in the same region [6] as well as in Nairobi [23] thus affirming subtype 16 as key player in CIN prevalence locally.

Analysis of risk determinants revealed that widows were over ten times more likely to test positive for CIN compared to the married counterpart. Studies have shown that being married is associated with early diagnosis of CIN and a more favorable prognosis for cervical cancer [27] possibly owing to partner support in regular clinic visit for screening and treatment. In this study, we




Fig. 2 Distribution rate of CIN among hr-HPV subgroups ($n = 95$)

Table 2 Correlates associated with CIN positivity by demographic and clinical characteristics

Characteristic	Univariate					Multivariable		
	N	Event N	OR ¹	95% CI ¹	p-value	OR ¹	95% CI ¹	p-value
Age category	189	95			0.180			
≤ 30 years			ref.	—		ref.	—	
> 30 years			1.81	0.76, 4.54		1.27	0.41, 4.21	0.700
Marital status	189	95			< 0.001			
Married			ref.	—		ref.	—	
Single			0.80	0.10–4.96		2.32	0.24, 19.6	0.400
Widow			11.4	3.18–73.2		11.9	2.53, 90.1	0.005
parity	189	95			0.460			
≤ 4 children			ref.	—				
> 4 children			1.27	0.68–2.38				
Contraceptives types	189	95			0.017			
None			ref.	—		ref.	—	
Hormonal-IUD			1.43	0.28–10.9		1.85	0.28–16.4	0.500
Implant Jadele/Norplant			0.40	0.15–0.96		1.18	0.37–3.86	0.800
injectable /Depo-provera			0.28	0.11–0.67		0.81	0.25–2.63	0.700
Oral pill			0.20	0.02–1.24		0.72	0.08–5.47	0.800
Contraceptive duration	189	95			0.700			
≤ 20			ref.	—				
> 20 years			1.16	0.52–2.63				
HIV status	189	95			< 0.001			
negative			ref.	—		ref.	—	
positive			2.79	1.56–5.10		2.43	1.25, 4.79	0.009
HSV2	189	95			0.024			
Negative			ref.	—		ref.	—	
Positive			2.41	1.12–5.46		2.70	1.15, 6.69	0.026
Chlamydia	189	95			< 0.001			
Negative			ref.	—		ref.	—	
Positive			3.83	1.84–8.51		3.63	1.61, 8.65	0.002

Regression analysis. ¹OR = Odds Ratio; CI = Confidence Interval; ref = reference; HSV-2, Herpes simplex virus-2; HPV, Human papilloma virus

found widowhood significantly associated with higher Odds of developing CIN. The protective effect of marriage on cervical cancer based on its sexually transmitted etiology have been reported widely showing that unmarried women would more often likely to have multiple sexual partners, inadequate access to health care and lack spousal emotional and social support essential for relieving psychosocial stress and unfavorable health outcomes [27]. Although studies have shown that older women are more vulnerable in terms of developing cervical cancer, especially high-stage disease [28]; in our study, we could not find any association between age and CIN. However, early onset of CIN among teenagers have been reported globally, possibly attributed to changes in diet, lifestyle, obesity, environment and the microbiome all of which interact with genomic and genetic susceptibilities [29] suggesting a possible aging dynamics as a risk factor.

Further analysis of risk factors revealed that women co-infected with HIV were twice more likely to test positive for CIN compared to their HIV-uninfected counterparts implying that, co-infection with HIV is independently and significantly associated with higher

Odds of developing CIN. Similar observation have been reported by Perez-Gonzalez *et al.* and Sosso *et al.*; and all noted that HPV infection was common among people living with HIV (PLWH) who were at greater risk of developing CIN [7, 30]. Indeed, PLWH have been shown to experience increased risk of persistent HPV infection including high viral load essential for CIN aetiology [31] more so when co-infection involve multiple HPV genotypes [23]. Few studies have explored the relationship between HSV-2 co-infection and the severity of cervical lesion. In our study, women co-infected with HSV-2 were twice more likely to test positive for CIN compared to their HSV-2-uninfected counterparts suggesting that co-infection with HSV-2 is significantly associated with higher Odds of developing CIN. It's good to note that HSV-2 co-infection has been incriminated with the initiation of oncogenic processes that are eventually picked by HPV in driving cervical cancer development [32]. Our finding corroborated a systematic review and a meta-analysis by Zhang *et al.* [33] showing that women co-infected with HSV-2 were 3 times more likely to test positive for CIN. Elsewhere, in another systematic

review, higher frequency of HSV-2 was observed among women experiencing invasive cervical cancer [32] suggesting a possible association that could potentially be attributed to shared immunological compromise, genetic predisposition, or lifestyle factors. In other studies, it has been suggested that genital HSV-2 infection possibly act in conjunction with HPV infection to increase modestly the risk of cervical intraepithelial neoplasia [34] of which investigation is still in progress. Locally, high burden of HSV-2 of 31.5% in young adults, 10.7% in adolescent youth [35] and 44% in the general population [36] have been reported in Kisumu and Kilifi, Kenya indicating the possibility of exposure to this potential risk factor.

Women co-infected with *Chlamydia trachomatis* (CT) were three times more likely to test positive for CIN compared to their CT-uninfected counterparts implying that co-infection with CT is equally associated with higher Odds of developing CIN. Again, this was in concurrence with a finding by Lu et al., showing prevalence of 21.8% low grade squamous intraepithelial lesions (LSIL) and 10.8% high grade squamous intraepithelial lesions (HSIL) among women co-infected with CT / HPV [37], and consistent with findings from neighborhood Uganda recording a significant association between CT / HPV co-infection and the development of LSIL [24]. Indeed, the DNA–host integration of HPV-CT is a useful event in the pathogenesis of HPV/CT related cancer the mechanism of which is still under investigation [14]. In a more related study, co-infection with CT was seen to support HPV persistence by suppressing the functions of Langerhans cells (LCs) pathways which are involved in the regulation of immune responses. Besides, the infection impairs LC functions by reducing the antigen-presenting ability and density of LCs; alter T-cell subsets, resulting in fewer CD4+ and CD8+ T cells and more infiltrating Tregs; decreases the CD4+/CD8+ T cell ratio to below 1; and induces greater T lymphocytes' apoptosis, hence impairing cell-mediated immunity and accelerating the progression of CIN [37]. In Rome Italy, women co-infected with CT / HPV were found to experience high frequency of high grade cervical lesions compared to their counterparts infected with HPV only [38], affirming the important role played by CT in cervical carcinogenesis. In this region, about 7.7% prevalence of CT have been reported among ANC mothers [39], with another 7.5% prevalence among women attending outpatient clinic [16], including 5.2% prevalence of anorectal CT and 12.8% prevalence of urethral CT among men having sex with men (MSM) [40], all in Kisumu, Kenya; thus supporting the need for surveillance of this potential risk factor for CIN.

Women on hormonal contraceptives, particularly hormone-containing intra uterine device (HIUD) were more likely to test positive for CIN compared to the none users'; suggesting that contraceptives are associated

with higher Odds of developing CIN. Contraceptives are mainly divided into progestin, estrogen or combined and their main actions are to inhibit ovulation, hinder the combination of sperm and eggs, and interfere with the implantation of fertilized eggs to achieve the purpose of contraception [41]. However, of late, several studies have linked the use of both oral contraceptives (OC) and HIUD with increased risk of developing CIN among the current and even recent long-term users of more than 5 years [41–44]. Use of HIUDs causes inflammatory and immunosuppressive changes that potentially affect the risk of persistent human papillomavirus infection and precancerous cervical lesions [44]. Ordinarily, the risk pattern among any hormonal and combined contraceptive users would increase with longer duration of use, but decline upon stopping, possibly taking longer to disappear among the prolonged users [45]. To the contrary, our study did not find association between OC use and CIN, neither did we find duration of use a risk factor. Elsewhere, use of HIUD was associated with productive and transient hr-HPV infections while OC was associated with transforming and integrated hr-HPV infections [42]. However, combined hormonal contraceptives are known to stimulate the integration of hr-HPV-DNA into the host genome, enhancing the expression of E6 and E7 HPV oncoproteins which stimulate the degradation of P53 tumor suppressor genes, increasing the ability of the viral DNA to transform cells leading to carcinogenesis [42].

To this end, it is good to note that previous local studies have mainly focused on the relationship between cervical cytology results and sexually transmitted pathogens majorly HIV and HPV [6, 23]. Conversely, cervical cytology results do not adequately represent the true picture of the cervix. Instead, cervical biopsy provides the gold standard for assessing cervical abnormalities, highlighting the need for assessing the relationship between sexually transmitted pathogens, HPV infection, and histological findings. In our study, we found a strong association between HIV, CT and HSV-2 co-infection with hr-HPV and CIN prevalence. Additionally, widowhood and use of HIUD was associated with CIN. This study had some limitations. First, the participants were recruited from a gynecological clinic having been referred from peripheral facilities with either vaginal or cervical abnormalities. This facility-based recruitment coupled with structural inefficiency in some rural setting that conveyed referrals limit the generalization of the study findings.

Conclusion

Co-infections with HIV, HSV-2 or *Chlamydia trachomatis* and use of HIUD were associated with increased risk of testing positive for CIN in HPV positive women in

western Kenya. Although the overall prevalence of CIN was high, high-grade CIN2 + was comparable to the rates reported earlier. Therefore, population screening for co-infections alongside hr-HPV is desirable and is likely to reduce the burden of CIN in the region. Besides, women positive for hr-HPV and opting for contraceptive ought to be counseled about the possible positive and negative side-effects of different contraception options.

Abbreviations

CI	confidence interval
SD	standard deviation
VL	viral load
CT	Chlamydia Trachomatis
CC	Cervical Cancer
CIN 1, 2, 3	Cervical Intraepithelial Neoplasia (Grade 1, Grade 2, Grade 3)
ICO-IARC	Catalan Institute of Oncology / International Agency for Research on Cancer
HPV	Human Papilloma Virus
HIV	Human Immunodeficiency virus
HR-HPV	High Risk Human Papilloma Virus
JOOTRH	Jaramogi Oginga Odinga Teaching and Referral Hospital

Acknowledgements

Appreciation to the department of health Kisumu county especially the staffs of JOOTRH for support in data collection. We equally appreciate Sustainable Development for Health on HIV (SD4H) program of Maseno University for supporting review of manuscript through organized workshop mentorship program; more so in statistical data analysis as provided by Prof. Lucas Othuon. Thanks to the Association of Kenya Medical Laboratory Scientific Officers (AKMLSO) for organised conference and workshop that equally supported data review. More thanks to Dr. Thomas Ongalo and Felix Humwa for additional support in data analysis.

Author contributions

CGO1, LO1, and BG1 conceived the research topic and design CGO1, and LO1 participated in acquisition, analysis and interpretation of data, CGO1, LO1, BG1, reviewed and approved the manuscript.

Funding

Research reported in this publication was supported in part by Fogarty International Center of the National Institute of Health (NIH) under Award Number D43TW011306. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Mental Health (NIMH). The NIMH had no role in the design and conduct of the study; collection, management, analysis, interpretation of data, review or approval of the manuscript; and decision to submit the manuscript for publication.

Data availability

Availability of data and materials. Data used in this study are included in the manuscript. Additional datasets are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

The institutional review board of JOOTRH provided ethics approval of the study number ERC.IB/VOL.1/602. Written informed consent was obtained from all participants.

Consent for publication

Consent was not needed as we used de-identified data.

Competing interests

The authors declare no competing interests.

Author details

¹Department of Biomedical Science and Technology, Maseno University, P.O Box Private Bag Maseno, Maseno 40105, Kenya

Received: 19 August 2024 / Accepted: 30 October 2024

Published online: 24 February 2025

References

1. ICO/IARC. Human Papillomavirus and Related Diseases Report [Internet]. 2023; <http://www.hpvcentre.net/>
2. Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin*. 2024;74:229–63.
3. Hassan MA, Itsura P, Odongo BE. Colposcopic and Histopathologic Comparative Interpretations Among Patients Undergoing Evaluation for Cervical Dysplasia in Western Kenya. *EMJ Repro Health* [Internet] 2024 [cited 2024 Aug 17]; <https://www.emjreviews.com/reproductive-health/article/colposcopic-and-histopathologic-comparative-interpretations-among-patients-undergoing-evaluation-for-cervical-dysplasia-in-western-kenya/>
4. Mwenda V, Mburu W, Bor JP, Nyangasi M, Arbyn M, Weyers S et al. Cervical cancer programme, Kenya, 2011–2020: lessons to guide elimination as a public health problem. *ecancer* [Internet] 2022 [cited 2024 Aug 14];16. <https://ecancer.org/en/journal/article/1442-cervical-cancer-programme-kenya-2011-2020-lessons-to-guide-elimination-as-a-public-health-problem>
5. Mungo C, Omoto J, Gwer S, Wameyo A, Kays M, Ganda G. Toward Cervical Cancer Elimination: evaluation of Access to Diagnostic services after Referral to a specialist gynecologist clinic at a Major Referral Hospital in Kisumu, Kenya. *JCO Global Oncol*. 2020;6:35–35.
6. Orango EO, Were E, Rode O, Muthoka K, Byczkowski M, Sartor H, et al. Novel concepts in cervical cancer screening: a comparison of VIA, HPV DNA test and p16INK4a/Ki-67 dual stain cytology in Western Kenya. *Infect Agents Cancer*. 2020;15:57.
7. Pérez-González A, Cachay E, Ocampo A, Poveda E. Update on the Epidemiological Features and clinical implications of human papillomavirus infection (HPV) and human immunodeficiency virus (HIV) Coinfection. *Microorganisms*. 2022;10:1047.
8. Banadakoppa Manjappa R, Bhattacharjee P, Shaw SY, Gitonga J, Kioko J, Songok F, Emmanuel F, Arimi P, Musyoki H, Masha RL, Blanchard J. A sub-national HIV epidemic appraisal in Kenya: a new approach for identifying priority geographies, populations and programmes for optimizing coverage for HIV prevention. *J Int AIDS Soc*. 2024;27:e26245.
9. García C, Hernández-García D, Valencia C, Rojo-León V, Pérez-Estrada JR, Werner M, et al. E6/E7 oncogenes in epithelial suprabasal layers and estradiol promote cervical growth and ear regeneration. *Oncogenesis*. 2017;6:e374–374.
10. Chachage M, Parikh AP, Mahenge A, Bahemana E, Mnkai J, Mbuya W, et al. High-risk human papillomavirus genotype distribution among women living with and at risk for HIV in Africa. *AIDS*. 2023;37:625–35.
11. Li M, Nyabigambo A, Navvuga P, Nuwamanya E, Nuwasiima A, Kaganda P, et al. Acceptability of cervical cancer screening using visual inspection among women attending a childhood immunization clinic in Uganda. *Papillomavirus Res*. 2017;4:17–21.
12. Yu X, He S. The interplay between human herpes simplex virus infection and the apoptosis and necroptosis cell death pathways. *Virol J*. 2016;13:77.
13. Macharia VM, Ngugi C, Lihana R, Ngayo MO, Transmitted. HIV-1 Drug resistance and the Role of Herpes Simplex Virus – 2 Coinfection among Fishermen along the Shores of Lake Victoria, Kisumu, Kenya. *J HIV Retrovirus* [Internet] 2016 [cited 2024 Aug 14];02. <http://hiv.imedpub.com/transmitted-hiv1-drug-resistance-and-the-role-of-herpes-simplex-virus-2-coinfection-among-fishermen-along-the-shores-of-lake-victo.php?aid=17345>
14. Arcia Franchini AP, Iskander B, Anwer F, Oliveri F, Fotios K, Panday P et al. The Role of Chlamydia Trachomatis in the Pathogenesis of Cervical Cancer. *Cureus* [Internet] 2022 [cited 2024 Aug 14]; <https://www.cureus.com/articles/70826-the-role-of-chlamydia-trachomatis-in-the-pathogenesis-of-cervical-cancer>
15. Mi Y, Gurumurthy RK, Zadora PK, Meyer TF, Chumduri C. *Chlamydia trachomatis* inhibits homologous recombination repair of DNA breaks by interfering with PP2A signaling. *Volume 9. mBio*; 2018. pp. e01465–18.

16. Nyakambi M, Waruru A, Oladokun A. Prevalence of genital Chlamydia trachomatis among women of reproductive age attending outpatient clinic at Kisumu County Referral Hospital, Kenya, 2021. *J Public Health Afr*. 2022;13:9.
17. Roura E, Travie N, Waterboer T, De Sanjosé S, Bosch FX, Pawlita M, et al. The influence of hormonal factors on the risk of developing Cervical Cancer and Pre-cancer: results from the EPIC Cohort. *PLoS ONE*. 2016;11:e0147029.
18. Gadducci A, Cosio S, Fruzzetti F. Estro-progestin contraceptives and risk of Cervical Cancer: a debated issue. *Anticancer Res*. 2020;40:5995–6002.
19. MoH. Kenya Demographic and Health Survey 2022. Key Indicators Report. 2023.
20. Kabanda R, Kiconco A, Ronald A, Beyer KMM, John SA. Correlates of intention to screen for cervical cancer among adult women in Kyotera District, Central Uganda: a community based cross-sectional study. *BMC Women's Health*. 2024;24:296.
21. Okyere J, Ayeberg C, Dosoo AK, Dickson KS. Cervical cancer screening among women with comorbidities: evidence from the 2022 Tanzania demographic and health survey. *BMC Public Health*. 2024;24:1093.
22. Khozaim K, Orango E, Christoffersen-Deb A, Itsura P, Oguda J, Muliro H, et al. Successes and challenges of establishing a cervical cancer screening and treatment program in western Kenya. *Int J Gynecol Obstet*. 2014;124:12–8.
23. Kangethe JM, Gichuhi S, Odari E, Pintye J, Mutai K, Abdullahi L et al. Confronting the human papillomavirus–HIV intersection: Cervical cytology implications for Kenyan women living with HIV. *South. Afr. j. HIV med.* [Internet] 2023 [cited 2024 Aug 14];24. <https://sajhivmed.org.za/index.php/hivmed/article/view/1508>
24. Sseddyabane F, Amnia DA, Mayanja R, Omonigbo A, Ssuuna C, Najjuma JN, et al. HPV-Chlamydial Coinfection, Prevalence, and Association with Cervical Intraepithelial lesions: a pilot study at Mbarara Regional Referral Hospital. *J Cancer Epidemiol*. 2019;2019:1–7.
25. Smith JS, Backes DM, Hudgens MG, Bailey RC, Veronesi G, Bogaarts M, et al. Prevalence and risk factors of human papillomavirus infection by penile site in uncircumcised Kenyan men. *Int J Cancer*. 2010;126:572–7.
26. Onyango CG, Ogonda L, Guyah B, Shiluli C, Ganda G, Orango OE, et al. Novel biomarkers with promising benefits for diagnosis of cervical neoplasia: a systematic review. *Infect Agents Cancer*. 2020;15:68.
27. El Ibrahimy S, Pinheiro PS. The effect of marriage on stage at diagnosis and survival in women with cervical cancer: marriage and cervical cancer stage and survival. *Psycho-oncology*. 2017;26:704–10.
28. Bønnekke S, Blaakær J, Steiniche T, Iachina M. Social factors and age play a significant role in cervical cancer and advanced-stage disease among Danish women. *BMC Cancer*. 2024;24:259.
29. Ugai T, Sasamoto N, Lee HY, Ando M, Song M, Tamimi RM, et al. Is early-onset cancer an emerging global epidemic? Current evidence and future implications. *Nat Rev Clin Oncol*. 2022;19:656–73.
30. Sosso SM, Tchouaket MCT, Fokam J, Simo RK, Semengue ENJ, Sando Z, et al. Human papillomavirus positivity and cervical lesions in relation to HIV infection: a comparative assessment in the Cameroonian female population. *J Public Health Afr*. 2023;14:7.
31. Menon S, Wusiman A, Boily MC, Kariisa M, Mabeya H, Luchters S, et al. Epidemiology of HPV genotypes among HIV positive women in Kenya: a systematic review and Meta-analysis. *PLoS ONE*. 2016;11:e0163965.
32. Sausen D, Shechter O, Gallo E, Dahari H, Borenstein R. Herpes Simplex Virus, Human Papillomavirus, and Cervical Cancer: overview, relationship, and treatment implications. *Cancers*. 2023;15:3692.
33. Zhang H, Cai S, Xia Y, Lin Y, Zhou G, Yu Y, et al. Association between human herpesvirus infection and cervical carcinoma: a systematic review and meta-analysis. *Viol J*. 2023;20:288.
34. Zhao Y, Cao X, Zheng Y, Tang J, Cai W, Wang H, et al. Relationship between cervical disease and infection with human papillomavirus types 16 and 18, and herpes simplex virus 1 and 2. *J Med Virol*. 2012;84:1920–7.
35. Akinyi B, Odhiambo C, Otieno F, Inzaule S, Oswago S, Kerubo E, et al. Prevalence, incidence and correlates of HSV-2 infection in an HIV incidence adolescent and adult cohort study in western Kenya. *PLoS ONE*. 2017;12:e0178907.
36. Nyiro JU, Sanders EJ, Ngetsa C, Wale S, Awuondo K, Bukusi E, et al. Seroprevalence, predictors and estimated incidence of maternal and neonatal herpes simplex virus type 2 infection in semi-urban women in Kilifi, Kenya. *BMC Infect Dis*. 2011;11:155.
37. Lu Y, Wu Q, Wang L, Ji L. Chlamydia trachomatis enhances HPV persistence through immune modulation. *BMC Infect Dis*. 2024;24:229.
38. Mancini F, Vescio F, Mochi S, Accardi L, di Bonito P, Ciervo A. HPV and Chlamydia trachomatis coinfection in women with pap smear abnormality: baseline data of the HPV Pathogen ISS study. *Le Infezioni in Medicina* 2018;2.
39. Mogaka JN, Drake AL, Matemo D, Kinuthia J, McClelland RS, Unger JA, et al. Prevalence and predictors of Chlamydia trachomatis and Neisseria gonorrhoeae among HIV-Negative pregnant women in Kenya. *Sex Trans Dis*. 2024;51:65–71.
40. Quilter LAS, Obondi E, Kunzweiler C, Okall D, Bailey RC, Djomand G, et al. Prevalence and correlates of and a risk score to identify asymptomatic anorectal gonorrhoea and chlamydia infection among men who have sex with men in Kisumu, Kenya. *Sex Transm Infect*. 2019;95:201–11.
41. Guo C, Zhan B, Li MY, Yue L, Zhang C. Association between oral contraceptives and cervical cancer: a retrospective case–control study based on the National Health and Nutrition Examination Survey. *Front Pharmacol*. 2024;15:1400667.
42. Loopik DL, Int'Hout J, Melchers WJG, Massuger LFAG, Bekkers RLM, Siebers AG. Oral contraceptive and intrauterine device use and the risk of cervical intraepithelial neoplasia grade III or worse: a population-based study. *Eur J Cancer*. 2020;124:102–9.
43. Abila DB, Wasukira SB, Ainembabazi P, Wabinga H. Burden of risk factors for cervical Cancer among women living in East Africa: an analysis of the latest demographic health surveys conducted between 2014 and 2017. *JCO Global Oncol* 2021;1116–28.
44. Skorstengaard M, Lynge E, Napolitano G, Blaakær J, Bor P. Risk of precancerous cervical lesions in women using a hormone-containing intrauterine device and other contraceptives: a register-based cohort study from Denmark. *Hum Reprod*. 2021;36:1796–807.
45. Iversen L, Fielding S, Lidegaard Ø, Hannaford PC. Contemporary hormonal contraception and cervical cancer in women of reproductive age. *Int J Cancer*. 2021;149:769–77.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.